

MEDICATIONS and SUPPLEMENTS which INCREASE BRUISING

If you are taking any of the following upon direction of a physician, consult that doctor before making any changes

Preparations Containing Aspirin or Salicylic Acid Derivatives should be avoided 3 weeks before surgery and 1 week after to minimize bruising

217	Buffaprin	Gemisyn
217 Strong	Buffasal	Genaced
4-Way Cold tablets	Bufferin	Genacote
Acetylsalicylic Acid	Buffets II	Genprin
Acuprin 81	Buffex	Gensan
Adult Analgesic pain reliever	Buffinol	Goody's extra strength
Aggrenox	Butalbital	Goody's Headache powder
Alka-Seltzer	C2	Halfprin
Anacin	CP-2	Headache tablet
Analval	Calmine	Healthprin
Anodynos	Cama arthritis pain reliever	Herbopyrine
Antidol	Carisoprodol compound	Instantine
Anthropan	Children's aspirin	Isollyl Improved
APAC Improved	Clinoril	Kalmex
APC	Congesprin chewables	Lanorinal
APO-ASA	Cope	Lortab with ASA
APO-ASEN	Coryphen Damason-P	Magnaprin
Arco Pain	Darvon compound	Marnal
Argesic	Darvon compound-65	Measurin
Arthritis	Darvon with ASA	Methcarbamol with aspirin
Arthritis Pain Formula	Darvon-N with ASA	Micrainin
Artria SR	Dasin	Meprobamate and aspirin
Arthrotec	Disalcid	Meprogesic Q
ASA	Dolcin	Midol
Ascriptin	Dolomine	Mobidin
Ascodeen	Dolprn #3 tablets	Mobigesic
Aspercin	Drinophen	Momentum
Aspergum	Duradyne	Neocylate
Aspermin	Easprin	Neogesic
Aspirin	Ecotrin	Nervine
Aspirin with codeine	Ecotrinol	Night-time Effervescent cold
Aspir-Low	Emagrin	Norgesic
AspirTab	Empirin	Norwich aspirin
Aspirtab Max	Empirin with codeine	Novasen
Astone	Emprazil	Orphenagesic
Astrin	Encaprin	Orphenagesic Forte
Atromid	Entrophen	Oxycodone and aspirin
Axotal	Equagesic	P-A-C
Azdone tablets	Equazine-M	Pain Aid
Azolid	Excedrin	Pain reliever tablets
B-A-C tablets	Fiogesic tablets	Pabrin buffered tabs
Baby aspirin	Fiorgen PF	Panodynes
Bayer aspirin	Fiorinal	Pepto-Bismol
Bayer children's cold tabs	Fiorinal with codeine	Percodan
BC powder or tablets	Gelpirin tablets	Percodan-Demi

Persistin	Saleto	Synalgos-DC capsules
Phenetron compound	Salflex	Talwin compound
Phenylbutaxone	Salicylamide	Tenol-Plus
PMS with ASA	Salicylate	Triaminicin
Presalin	Salicylic Acid	Trigesic
Propoxyphene compound	Salocol	Trilisate
Propoxyphene napsylate	Salsalate	Tri-Pain
Quiet World tablets	Sine-Off sinus medicine	Ursinus Inlay-Tabs
Rhinoceps	Sloprin	Valesin
Robaxisal	Soma compound	Vanquish
Roxiprin	Soma compound with codeine	Verin
S-A-C	St. Joseph	Wesprin
Salabuff	Stanback powder	Zorprin
Salatin	Supac	

**Preparations Containing Mostly Non-steroidal Anti-inflammatory Agents
should be avoided 1-2 weeks before injections to minimize the chance of bruising**

Aches-N-Pain	Diclofenac	Magsal	Nu-Pirox
Actiprofen	Diffunisal	Marnal	Nuprin
Actron	Dimetapp Sinus	Measurin	Nu-Sulindac
Addaprin	Dolgesic	Meclofenamate	Oraflex
Advil	Dolobid	Meclomen	Orudis
Aleve	Dristan Sinus	Medipren	Oruvail
Anaprox	Duoprin	Mefenamic	Oxaprozin
APC	Durasal	Meloxicam	Pabalate-SF tabs
Ansaid	EC-Naprosyn	Menadol	Pamprin-IB
APO-Diclo	Etodolac	Midol	PediaProfen
APO-Diflunisal	Excedrin IB	Mobic	Phenylbutazone
APO-Flurbiprofen	Feldene	Motrin	Piroxicam
APO-Ibuprofen	Fenoprofen	Nalfon	Ponstan
APO-Indomethacin	Four-way cold tabs	Naprelan	Ponstel
APO-Keto	Flurbiprofen	Napron X	Q-Profen
APO-Napro	Froben	Naprosyn	Relafen
APO-Naproxen	Gaysal	Naproxen	Rhodis
APO-Piroxicam	Genpril	Naxen	Rufen
APO-Sulin	Haltran	Novo-Difenac	Saleto
Arthrotec	Ibifon 600	Novo-Diflunisal	Sine-Aid IB
Atromid	Ibren	Novo-Flurprofen	Sulindac
Azolid	IBU	Novo-Keto-EC	Synflex
Bayer Select	Ibuprin	Novo-Methacin	Tolectin
Buf-tabs	Ibuprofen	Novo-Naprox	Tolmetin
Butazolidin	Ibuprohm	Novo-Pirocam	Toradol
Cataflam	Indochron ER	Novo-Profen	Trendar
Clinoril	Indocid	Novo-Sundac	Ultraprin
Clofibrate	Indocin	Novo-Tolmetin	Unipro
CoAdvil	Indomethacin	Nu-Diclo	Valprin
Cotybutazone	Ketoprofen	Nu-Flubiprofen	Voltaren
Cramp End	Ketorolac	Nu-Ibuprofen	Voltaren
CP-2 tabs	Lodine	Nu-Indo	
Daypro	Magan	Nu-Naprox	

Prescription Medications That Increase Bruising

Baclofen	Duragesic patch	Parnate	Ticlid
Cilostazol	Etrafon	Perfenazine	Ticlopidine
Clopidogrel	Fentanyl	Persantine	Tilafon
Coumadin	Lioresal	Plavix	Warfarin
Dipridacot	Panalgesic	Pletal	
Dipyridamole	Pan Warfarin	Selenium	

Herbal Supplements That Increase Bruising

Bilberry	Fish oil	Soma
Chamomile	Flaxseed	St. John's Wort
Dong quai root	Garlic	Valerian
Echinacea	Ginger	Vitamin E
Ephedra	Ginkgo	Willow bark
E vitamin	Ginseng	Yohimbe
Feverfew	Kava	

IMPORTANT

If you are taking any of the previous medications or dietary supplements upon recommendation or direction of a physician, consult that doctor before making any changes